

SUPPLEMENTAL LABELING

This supplemental label expires on 01/08/2023 and must not be used or distributed after this date.

CLETHODIM GROUP 1 HERBICIDE



Supplemental Label to Add Camelina.

ACTIVE INGREDIENT:	(% by weight)
Clethodim: (E)-2-[1-[(3-chloro-2-propenyl)oxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one	26.4%
OTHER INGREDIENTS:	73.6%
TOTAL:	100.0%
Contains petroleum distillate.	
Contains 2.0 lbs. clethodim per gallon.	

**KEEP OUT OF REACH OF CHILDREN
WARNING—AVISO**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the label affixed to the container for **Ceridian 2 EC** before applying.

This labeling must be in possession of the user at the time of application.

Use of **Ceridian 2 EC** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Ceridian 2 EC**.

EPA Reg. No.: 91234-154
20200109a

CROP SPECIFIC RESTRICTIONS AND LIMITATIONS

Crops ¹	Minimum Time from Application to Harvest (PHI)	Use Rate Per Acre	Crop Oil Concentrate Rates Per Acre ²	Special Use Instructions
<p>Canola including Rapeseed Subgroup 20A* except flax seed, mustard seed and sesame seed</p> <p>Borage Crambe Cuphea Echium Echium Gold of Pleasure (Camelina) Hare's Ear Mustard Lesquerella Lunaria Meadowfoam Milkweed Oil Radish Poppy seed Rapeseed (canola) Sweet Rocket</p>	<p align="center">70 days</p>	<p align="center">4-6 fl. oz. (0.0625 – 0.09375 lb a.i)</p>	<p align="center">1.0% v/v in the finished spray volume</p>	<p>Verify crop safety to Ceridian 2 EC on a small area of the crop, at the desired Ceridian 2 EC rate and with the same non-ionic surfactant (NIS) that will be used on the field. If no crop response is evident 7 days after treatment, Ceridian 2 EC may be used on the entire field at the rate tested and with the same NIS used in the crop safety test.</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 6 fl. oz./A (0.09375 lb a.i./A) in a single application. • DO NOT make more than 1 application per acre per year. • DO NOT apply more than 6 fl. oz./A (0.09375 lb a.i./A) in a year. • DO NOT apply after crop has begun bolting. Crop injury may occur when this product is applied during the bloom period <p>*Not for Use in California</p>

¹ **Ceridian 2 EC** is not to be used on vegetable crops being grown for seed production unless specific use directions are provided.

Manufactured for:
Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513